

# Generalization of Weierstrassian Elliptic Functions to $\mathbf{R}^n$

Cihan Saçlıoğlu<sup>1</sup>

TÜBİTAK - Marmara Research Center  
Research Institute for Basic Sciences  
Department of Physics  
41470 Gebze, Turkey

February 1, 2008

## Abstract

The Weierstrassian  $\wp, \zeta$  and  $\sigma$  functions are generalized to  $\mathbf{R}^n$ . The  $n = 3$  and  $n = 4$  cases have already been used in instanton solutions which may be interpreted as explicit realizations of spacetime foam and the monopole condensate, respectively. The new functions satisfy higher dimensional versions of the periodicity properties and Legendre's relations obeyed by their familiar complex counterparts. For  $n = 4$ , the construction reproduces functions found earlier by Fueter using quaternionic methods. Integrating over lattice points along all directions but two, one recovers the original Weierstrassian elliptic functions.

---

<sup>1</sup>Permanent address: Boğaziçi University, Department of Physics, 80626 Istanbul, Turkey.

Kaluza-Klein, Supergravity, String and Superstring theories and Super p-branes all involve dimensions beyond four, which then have to be compactified. The compactification results in lattices in the higher dimensions [1]. It might therefore be of interest to consider generalizations of doubly periodic functions to arbitrary  $\mathbf{R}^n$ . The  $\mathbf{R}^4$  case has already been treated by Fueter [2], who succeeded in obtaining quaternionic analogues of Weierstrassian elliptic functions. In this note, we will present a unique and straightforward extension of Fueter's results to  $\mathbf{R}^n$ . This general construction yields Fueter's functions in  $n = 4$  without requiring the use of quaternions. Furthermore, when one takes infinitesimal lattice spacings in all but two lattice directions and integrates over these points one recovers the usual Weierstrassian functions.

Apart from possible future physics applications of n-tuply periodic functions, it is worth noting that (quasi)periodic classical solutions of Gravity and Yang-Mills theory have already been considered. Thus, inspired by Rossi's observation [3] that a singly-periodic instanton configuration represents a BPS monopole [4], Gursev and Tze [5] used Fueter's hyperelliptic functions to construct a solution with one Yang-Mills instanton per spacetime cell. This solution may be related to the monopole condensate [6] or to the Copenhagen vacuum [7], which is also based on a periodic arrangement of the theory's solitons. Hawking, on the other hand, argued [8] that metrics with a single gravitational instanton per unit cell make the dominant contribution to the path integral of Einstein's gravity. An adaptation of Weierstrassian  $\sigma$  and  $\zeta$  functions to three dimensions [9] along the lines presented here leads to just such a self-dual solution, providing an explicit example of a "spacetime foam" based on Gibbons-Hawking multicenter metrics [10].

We start by examining how the infinite sums defining the Weierstrassian functions are made to converge. These are

$$\wp(z) = \frac{1}{z^2} + \sum_{\omega \neq 0} \left\{ \frac{1}{(z - \omega)^2} - \frac{1}{\omega^2} \right\}, \quad (1)$$

$$\zeta(z) = - \int \wp(z) dz = \frac{1}{z} + \sum_{\omega \neq 0} \left\{ \frac{1}{(z - \omega)} + \frac{1}{\omega} + \frac{z}{\omega^2} \right\}, \quad (2)$$

$$\rho(z) \equiv \ln \sigma(z) = \int dz \zeta(z) = \ln(z) + \sum_{\omega \neq 0} \left\{ \ln(z - \omega) - \ln(-\omega) + \frac{z}{\omega} + \frac{z^2}{2\omega^2} \right\}, \quad (3)$$

or,

$$\rho(z) = \ln(z) + \sum_{\omega \neq 0} \left\{ \ln\left(1 - \frac{z}{\omega}\right) + \frac{z}{\omega} + \frac{z^2}{2\omega^2} \right\} \quad (4)$$

In the above  $\omega = n_1\omega_1 + n_2\omega_2$ , with  $(\omega_1, \omega_2)$  defining the basic lattice cell  $\mathcal{C}$ . The parentheses  $\{\}$  ensure that each term in the sum is absolutely convergent; the series become meaningless if the parentheses are broken up and the terms they contain are separately summed. An efficient method for studying the convergence properties of (1)-(3) is provided by an integral test in which the double sum is replaced by  $\int_{|\omega|_{min}}^{\infty} d|\omega| |\omega|$ . This reveals why one, two and three subtraction terms are needed in (1), (2) and (3), respectively. Employing a simple dimensional argument, one can proceed further and determine the precise forms of the subtraction terms, given only the first term in the sum (3), for example. To do this, we note (a)  $\rho$  is dimensionless (assigning the dimension of length to the coordinate  $z$ , say); (b) in the power series expansion of (4), the highest power of  $\omega$  permitted by convergence requirements is  $\omega^{-3}$ . One then realizes that the subtraction terms following  $\ln(1 - \frac{z}{\omega})$  must be chosen so that  $\rho(z)$  has the expansion

$$\rho(z) = \ln z + \sum_{\omega \neq 0} O\left(\frac{z^3}{\omega^3}\right) + \dots \quad (5)$$

Hence using the power series

$$\ln(1 - z/\omega) = -z/\omega - z^2/2\omega^2 - z^3/3\omega^3 - \dots \quad (6)$$

one chooses the three subtraction terms

$$-\ln(-\omega) + z/\omega + z^2/2\omega^2 \quad (7)$$

to arrive at the form (5). This will be the key in constructing  $\mathbf{R}^n$  analogues of  $\rho(z)$ .

We will later need the well known facts that  $\wp(z)$  is doubly periodic while  $\zeta(z)$  and  $\sigma(z)$  are quasiperiodic with the transformation properties

$$\zeta(z + \omega_{1,2}) = \zeta(z) + \eta_{1,2} \quad (8)$$

and

$$\sigma(z + \omega_{1,2}) = -\sigma(z) \exp \eta_{1,2} (z + \omega_{1,2}/2) \quad (9)$$

where  $\eta_{1,2} = 2\zeta(\omega_{1,2}/2)$ .

We also record the so-called Legendre's relations

$$\oint_{\partial\mathcal{C}} \zeta(z) dz = 2\pi i \quad (10)$$

and

$$\oint_{\partial\mathcal{C}} \zeta(z) dz = 2\pi i = \eta_1\omega_2 - \eta_2\omega_1. \quad (11)$$

We first rewrite (10) in the rather unconventional form reminiscent of Gauss's theorem

$$\int_{\mathcal{C}} dV \nabla^2 \rho(z) = \oint_{\partial\mathcal{C}} d\vec{\sigma} \cdot \vec{\nabla} \rho(z) = 2\pi, \quad (12)$$

where  $dV = dxdy$  and the "surface element"  $d\vec{\sigma} = \hat{n}dl$ . Here  $dl$  is the arclength and  $\hat{n}$  a unit normal vector pointing outwards on  $\partial\mathcal{C}$ .

Let us next introduce the lattice basis vectors  $q_{\mu}^{(a)}$ , where  $a, \mu = 1, \dots, n$ . The volume of the unit cell  $\mathcal{C}$  is given by

$$V_n = \frac{1}{n!} \epsilon_{a_1 \dots a_n} \epsilon^{\mu_1 \dots \mu_n} q_{\mu_1}^{(a_1)} \dots q_{\mu_n}^{(a_n)}. \quad (13)$$

The basis vectors of the reciprocal lattice obey

$$r_{\mu}^{(a)} q_{\mu}^{(b)} \equiv r^{(a)} \cdot q^{(b)} = \delta^{ab}. \quad (14)$$

They are obtained from the  $q_{\mu}^{(a)}$  via

$$r_{\mu_1}^{(a_1)} = \frac{1}{(n-1)!V_n} \epsilon_{a_1 \dots a_n} \epsilon^{\mu_1 \dots \mu_n} q_{\mu_2}^{(a_2)} \dots q_{\mu_n}^{(a_n)}. \quad (15)$$

We now seek higher dimensional versions of (12) in the form

$$\int_{\mathcal{C}} dV_n \partial_{\mu} \partial_{\mu} \rho_n(x) = \oint_{\partial\mathcal{C}} d\sigma_{\mu} \partial_{\mu} \rho_n(x) = - \int d\omega_n, \quad (16)$$

where of course

$$\int d\omega_n \equiv \Omega_n = 2\pi^{n/2}/\Gamma(n/2). \quad (17)$$

In order to obtain the result (16), the function  $\rho_n(x)$ , which is to serve as the analog of  $\rho(z)$ , should have the form

$$\rho_n(x) \propto G_n(x) + \sum_{q \neq 0} \dots \sum G_n(x - q) + (\mathbf{R}^n \text{ harmonics}). \quad (18)$$

In (18),  $G_n(x)$  is the Green's function for  $\mathbf{R}^n$  obeying

$$\partial_\mu \partial_\mu G_n(x) = -\Omega_n \delta(x_1) \dots \delta(x_n) \equiv -\Omega_n \delta(x) \quad (19)$$

and

$$q = n_1 q^{(1)} + \dots + n_n q^{(n)}. \quad (20)$$

We have, of course

$$G_n(x - q) = \frac{1}{(x^2 - 2x \cdot q + q^2)^{\frac{n-2}{2}}} = \frac{1}{|x - q|^{n-2}} \quad (n > 2). \quad (21)$$

The harmonics in (18) should now be chosen according to the general strategy outlined between equations (4) and (8). Thus in order to render convergent the sum (18), whose integral counterpart contains terms behaving like  $\int_{|q|_{min}}^\infty d|q| |q|^{n-1}/|q|^{n-2}$ , one again needs 3 subtraction terms. These are in fact nothing but the first three terms in the MacLauren expansion of (21) for  $|x| \ll |q|_{min}$ . This immediately leads to

$$\rho_n(x) = \frac{1}{(x^2)^{\frac{n-2}{2}}} + \sum_{q \neq 0} \left\{ \frac{1}{|x - q|^{n-2}} - \frac{1}{|q|^{n-2}} \left[ 1 + \frac{(n-2)}{q^2} (q \cdot x + \frac{1}{2q^2} (n(q \cdot x)^2 - q^2 x^2)) \right] \right\} \quad (22)$$

Again, absolute convergence is attained only by considering each term defined by the outermost parentheses as an indivisible unit. Hence the terms proportional to  $q \cdot x$  and to  $1/2q^2$  cannot be summed separately (in which case they would appear to give zero by symmetry!) anymore than the  $\omega^{-1}$  term in (2) can. Note that the subtraction terms are indeed harmonics as anticipated in (18).

This is perhaps an appropriate point for comparing our results with Fueter's. Fueter introduces the unit quaternions  $\mathbf{e}_\mu$  and  $\bar{\mathbf{e}}_\mu$  corresponding to  $(I, \mathbf{e}_i)$  and  $(I, -\mathbf{e}_i)$ , respectively. We have, as usual,  $\mathbf{e}_i \mathbf{e}_j = -\delta_{ij} + \epsilon_{ijk} \mathbf{e}_k$ , the indices  $i, j, k$  running from 1 to 3. Using these, one defines  $\mathbf{x} = x_\mu \mathbf{e}_\mu$ ,  $\bar{\mathbf{x}} = x_\mu \bar{\mathbf{e}}_\mu$ ,  $D = \mathbf{e}_\mu \partial_\mu$  and  $\bar{D} = \bar{\mathbf{e}}_\mu \partial_\mu$ . One then has  $x^2 = \bar{\mathbf{x}} \mathbf{x}$  and  $\bar{D} D = \partial_\mu \partial_\mu$ . Then, starting with the function

$$\mathbf{Z}(\mathbf{x}) = \frac{1}{\mathbf{x}} + \sum_{q \neq 0} \left\{ \frac{1}{\mathbf{x} - \mathbf{q}} + \frac{1}{\mathbf{q}} + \frac{1}{\mathbf{q}} \mathbf{x} \frac{1}{\mathbf{q}} + \frac{1}{\mathbf{q}} \mathbf{x} \frac{1}{\mathbf{q}} \mathbf{x} \frac{1}{\mathbf{q}} + \frac{1}{\mathbf{q}} \mathbf{x} \frac{1}{\mathbf{q}} \mathbf{x} \frac{1}{\mathbf{q}} \mathbf{x} \frac{1}{\mathbf{q}} \right\}, \quad (23)$$

where  $\mathbf{q} = n_a \mathbf{q}^{(a)} = n_a q_\mu^{(a)} \mathbf{e}_\mu$ , the  $q^{(a)}$  being lattice vectors, one arrives at the quaternionic version of Weierstrass's  $\zeta(z)$  via

$$\zeta^{\mathbf{F}}(\mathbf{x}) = \partial_\mu \partial_\mu \mathbf{Z}(\mathbf{x}). \quad (24)$$

We will see later that (24) indeed transforms the same way as  $\zeta(z)$  under lattice shifts. The function corresponding to  $\ln \sigma(z)$  is

$$\rho(\mathbf{x}) = D\mathbf{Z}(\mathbf{x}) \quad (25)$$

and, remarkably, it turns out to have the quaternion-free form

$$\rho(x) = \frac{1}{x^2} + \sum_{q \neq 0} \left\{ \frac{1}{(x-q)^2} - \frac{1}{q^2} - \frac{2x \cdot q}{q^4} - \frac{1}{q^6} (4(q \cdot x)^2 - q^2 x^2) \right\}, \quad (26)$$

coinciding with the  $n=4$  case of (22).

Returning next to (12), we see that the analog of  $\zeta(z)$  is the  $n$ -gradient

$$\zeta_\mu^{(n)}(x) = \partial_\mu \rho_n(x). \quad (27)$$

Taking  $n=4$  and contracting both sides with the quaternion units  $\bar{\mathbf{e}}_\mu$ , we recover the  $\zeta^{\mathbf{F}}$  function of Fueter [2]

$$\zeta^{\mathbf{F}} = \bar{\mathbf{e}}_\mu \partial_\mu \rho_4 = \overline{D} \rho_4. \quad (28)$$

Leaving the quaternionic special case aside, let us now find the higher dimensional analogs of (8) and (9). Since

$$\partial_\mu \partial_\mu \rho_n = \partial_\mu \zeta_\mu^{(n)} = -\Omega_n \sum \dots \sum \delta(x-q) \quad (29)$$

is a perfectly  $n$ -tuply periodic distribution, a shift by a lattice basis vector can only change  $\zeta_\mu^{(n)}$  by a constant vector. Thus

$$\zeta_\mu^{(n)}(x + q^{(a)}) = \zeta_\mu^{(n)}(x) + \eta_\mu^{(n)(a)}. \quad (30)$$

This is indeed the transformation law for  $\zeta^{\mathbf{F}}$  when  $n=4$ . Putting  $x = -q^{(a)}/2$  and noting that  $\zeta_\mu^{(n)}(-x) = -\zeta_\mu^{(n)}(x)$ , we find

$$2\zeta_\mu^{(n)}(q^{(a)}/2) = \eta_\mu^{(n)(a)}. \quad (31)$$

Integrating (30) and using  $\rho_n(-x) = \rho_n(x)$ , we obtain

$$\rho_n(x + q^{(a)}) = \rho_n(x) + \eta^{(n)(a)} \cdot (x + q^{(a)}/2), \quad (32)$$

where there is no sum on the index  $(a)$  on the right hand side. Using (30), we can also obtain the higher dimensional form of Legendre's second relation

(11). Let us evaluate  $\oint_{\partial\mathcal{C}} d\sigma_\mu \zeta_\mu^{(n)}(x)$ , where  $\partial\mathcal{C}$  is the surface of the fundamental hyperparallelepiped. Noting that the normal to the hyperplane defined by  $\{q^{(a_2)}, \dots, q^{(a_n)}\}$  is along the reciprocal vector  $r^{(a_1)}$  given in (15), we find

$$V_n \sum_{a=1}^n \eta^{(n)(a)} \cdot r^{(a)} = \Omega_n = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \quad (33)$$

as the  $n$ -dimensional generalization of Legendre's second relation.

There is an interesting relationship between  $\rho_n$  and  $\rho_{n-1}$  which can be iterated until one finally arrives at the Weierstrassian elliptic functions. It is simplest to describe this relationship in the case of a rectangular lattice although it is generally valid. Starting with  $\rho_n$ , one first chooses one lattice direction to coincide with, say, the  $x_n$  axis and then lets the lattice spacing (in the same direction only) become infinitesimal. Then, integrating over  $q_n$ , one obtains  $\rho_{n-1}$  up to a multiplicative constant. Repeating this procedure until  $\rho_3$  is found, one finally has

$$-\int_{-\infty}^{\infty} dq_3 \rho_3 = \rho(z) + \rho(\bar{z}) = \int dz \zeta(z) + \int d\bar{z} \zeta(\bar{z}), \quad (34)$$

where the remaining lattice basis vectors  $(\vec{q}^{(1)}, \vec{q}^{(2)})$  provide the periods  $(\omega_1, \omega_2)$ . Thus the  $\rho_n$  presented here are indeed intimately connected with Weierstrassian functions. We also observe that the split into an analytic and anti-analytic part is a feature of two dimensions, not shared for example by Fueter's quaternionic functions.

Finally, the reader may wonder how the  $\mathbf{R}^n$  counterpart of  $\wp(z)$  is to be defined. It is clear that two derivatives of  $\rho_n(x)$  will be involved. It is natural to classify the resulting functions according to their  $SO(n)$  transformation properties. One has the symmetric traceless second rank  $SO(n)$  tensor

$$\wp_{\mu\nu}^{(n)} \equiv (\partial_\mu \partial_\nu - \frac{\delta_{\mu\nu}}{n} \partial_\lambda \partial_\lambda) \rho_n(x) \quad (35)$$

and the  $SO(n)$  scalar

$$\pi^{(n)} \equiv \partial_\lambda \partial_\lambda \rho_n = -\Omega_n \sum \dots \sum \delta(x - q) \quad (36)$$

already encountered in (29). Although both (35) and (36) are fully  $n$ -tuply periodic, the former involves true functions while the latter is an  $n$ -fold sum

over distributions.  $\wp_{\mu\nu}^{(n)}$  is also similar to  $\wp(z)$  in that

$$\oint_{\partial\mathcal{C}} d\sigma_\mu \wp_{\mu\nu}^{(n)} = \int_{\mathcal{C}} dV_n \partial_\mu \wp_{\mu\nu}^{(n)} = 0 \quad (37)$$

just like  $\wp(z)$  which obeys

$$\oint_{\partial\mathcal{C}} \wp(z) dz = 0. \quad (38)$$

I am grateful to A. Aliyev, O. F. Dayi, C. Delale and Y. Nutku for patiently instructing me in the use of LATEX and to my colleagues at the MRC for their hospitality.



## References

- [1] W. Lerche, A. N. Schellekens and N. P. Warner, Phys. Rep. **C 177** (1989) 1.
- [2] R. Fueter, C. R. Congress Internat. Math. Oslo 1 (1936) 75; Comment. Math. Helv. **7** (1934-1935) 307; Monatsh. Math. Phys. **48** (1939) 345.
- [3] R. Rossi, Nucl. Phys. **B 149** (1979) 170.
- [4] B. Bogomol'nyi, Sov. J. Nucl. Phys. **24** (1976) 860; M. K. Prasad and C. M. Sommerfield, Phys. Rev. Lett. **35** (1975) 760.
- [5] F. Gursey and H. C. Tze, Annals of Physics **128** (1980) 29.
- [6] S. Mandelstam, Phys. Rep. **C 67** (1980) 109.
- [7] H. B. Nielsen *in* Particle Physics 1980, ed. by I. Andric, I. Dadić and N. Zovko (North-Holland 1981) and references therein; J. Ambjorn and P. Olesen, Nucl. Phys. **B 170** (1980) 60.
- [8] S. W. Hawking *in* General Relativity, An Einstein Centenary Survey, ed. by S. W. Hawking and W. Israel (Cambridge University Press, 1970).
- [9] S. Nergiz and C. Saclioglu, "A Quasiperiodic Gibbons-Hawking Metric and Spacetime Foam", MRC-PH-TH-4-95, hep-th/950541.
- [10] G. W. Gibbons and S. W. Hawking, Phys. Lett. **78 B** (1978) 430.